

Sem-III - Thermal Physics

(Instructor: AKB, Department of Physics, Asutosh College)

Assignment II: 1st – 2nd law of Thermodynamics & Pure Substances

Submission due date: xx/xx/202x

Q.1) If a gas is both ideal and paramagnetic obeying Curie's law, show that the entropy is given by

$$S = c_{V,M} \ln T + nR \ln V - \frac{M^2}{2C'_c} + \text{constant} ,$$

where $c_{V,M}$ is the heat capacity at constant volume, magnetization assumed constant and C'_c is Curie's constant.

Q.2) A liquid is irregularly stirred in a well-insulated container and thereby undergoes a rise in temperature. If we regard the liquid as the system, **(a)** Has heat been transferred? **(b)** Has work been done? **(c)** What is the sign of ΔU ?

Q.3) The equation of state of a novel matter is $PV = AT^3$ with A a constant. The internal energy of the matter is $U = BT^n \ln(V/V_0) + f(T)$. Using first law of thermodynamics, find B and n .

Q.4) Suppose an engine works between two reservoirs at T_1 and T_2 ($T_2 > T_1$) until both reservoirs attain final temperature T_c . Show that $T_c > \sqrt{T_1 T_2}$. What is the maximum amount of work obtainable from this engine?

Q.5) A Carnot engine has an efficiency of 30% when the sink temperature is $27^\circ C$. What must be the change in temperature of the source to make its efficiency 50%?

Q.6) An inventor claims to have developed an engine working between $600K$ and $300K$ to deliver an efficiency of 52%. Using Carnot's theorem, can you decipher whether this claim is valid?

Q.7) Two Carnot engines X and Y are operating in series. X receives heat at $1200K$ and rejects to a reservoir at temperature TK . The second engine Y receives the heat rejected by X and in turn rejects to a heat reservoir at $300K$. Calculate the temperature T for the situation when, (i) The work output of two engines are equal, (ii) The efficiency of two engines are equal.

Q.8) A Carnot's refrigerator takes heat from water at $0^\circ C$ and discards it to a room temperature. $1Kg$ of water at $0^\circ C$ is to be changed into ice at $0^\circ C$. How many calories of heat are discarded to the room? What is the work done by the refrigerator in this process? What is the coefficient of performance [$P = Q_{cold}/(Q_{hot} - Q_{cold})$] of the machine? Given, room temperature is $27^\circ C$ and $1Cal = 4.2Joule$.

Q.9) A thermally conducting bar of length L , area A , density ρ is brought to a nonuniform temperature distribution by sandwiching between hot (temperature T_h) and cold reservoir (temperature T_c). The bar is removed from reservoirs, thermally insulated and kept at constant pressure. Show that the change in entropy of the bar is

$$\Delta S = c_p \rho A L \left\{ 1 + \ln \left(\frac{T_h + T_c}{2} \right) + \frac{T_c}{T_h - T_c} \ln T_c - \frac{T_h}{T_h - T_c} \ln T_h \right\}.$$

Q.10) Consider a metal (say Copper) at $300K$ with the following values, $V = 7.06 \text{ cm}^3/\text{mol}$, $K_T = 7.78 \times 10^{-12} \text{ N/m}^2$, $\beta = 50.4 \times 10^{-6} \text{ K}^{-1}$, $C_p = 24.5 \text{ J/molK}$. Determine C_v .

Q.11) Prove that the ratio of adiabatic $\left[\alpha_S = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_S \right]$ to isobaric $\left[\alpha_P = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_P \right]$ coefficient of expansion is $\frac{1}{1-\gamma}$. Also, prove that the ratio of adiabatic $\left[E_S = -V \left(\frac{\partial P}{\partial V} \right)_S \right]$ to isothermal $\left[E_T = -V \left(\frac{\partial P}{\partial V} \right)_T \right]$ elasticities is equal to the ratio of specific heats.

Q.12) Prove that the ratio of adiabatic $\left[\beta_S = \frac{1}{P} \left(\frac{\partial P}{\partial T} \right)_S \right]$ to isochoric $\left[\beta_V = \frac{1}{P} \left(\frac{\partial P}{\partial T} \right)_V \right]$ pressure coefficient of expansion is $\frac{\gamma}{\gamma-1}$.

Q.13) (a) If equation of state of certain material satisfies $P = \frac{RT}{V} \left(1 + \frac{B''}{V} \right)$ where $B'' = B''(T)$, show that

$$C_V = -\frac{RT}{V} \frac{d^2}{dT^2} (B''T) + C_V^\infty,$$

where C_V^∞ represents the value of C_V when V is very large. **(b)** In case $P = \frac{RT}{V} (1 + B'P)$ where $B' = B'(T)$, show that

$$C_P = RTP \frac{d^2}{dT^2} (B'T) + C_P^0,$$

where C_P^0 represents the value of C_P when pressure tends to zero.

Q.14) Using Berthelot's equation of state $P = \frac{RT}{V-b} - \frac{a}{TV^2}$, show that the critical constants are

$$P_c = \frac{1}{12b} \sqrt{\frac{2aR}{3b}}, \quad V_c = 3b, \quad T_c = \sqrt{\frac{8a}{27bR}}; \quad \frac{RT_c}{P_c V_c} = \frac{8}{3}.$$

Q.15) The boiling point of a liquid at pressure P_0 is T_0 . Its molar latent heat of vaporisation is L and molar volume of the liquid phase is negligible as compared to vapour phase. The vapour phase obeys the ideal gas equation. Show that the boiling point T at pressure P is given by,

$$\ln \left(\frac{P}{P_0} \right) = \frac{L}{RT_0} \left(1 - \frac{T_0}{T} \right).$$